arXiv:1304.4345v1 [hep-lat] 16 Apr 2013

Conformal Behavior in High Temperature QCD

K.-I. Ishikawa,[1] Y. Iwasaki,[2] Yu Nakayama,[3] and T. Yoshie[2]
[1] *Graduate School of Science, Hiroshima University,Higashi-Hiroshima, Hiroshima 739-8526, Japan*
[2] *Center for Computational Sciences, University of Tsukuba,Tsukuba, Ibaraki 305-8577, Japan*
[3] *California Institute of Technology, Pasadena, CA 91125, USA*

(Dated: April 17, 2013)

We give a new perspective on the properties of quarks and gluons at finite temperature T in $N_f = 2 \sim 6$ QCD. We point out the existence of an IR fixed point for the gauge coupling constant at $T > T_c$ (T_c is the chiral phase transition temperature). Based on this observation we predict theoretically and verify numerically that the correlation functions of a meson $G_H(t)$ at $T/T_c > 1$ decay with a power-law corrected Yukawa-type decaying form, $G_H(t) = c\exp(-m_H t)/t^\alpha$ in the "conformal region" defined by $m_H < c\Lambda_{\rm IR}$, where $\Lambda_{\rm IR}$ is the IR cutoff, m_H is the ground state energy in the cannel H and c is a constant of order 1. The decaying form is the characteristics of conformal theories with an IR cutoff. We discuss in detail how the resulting hyperscaling relation of physical observables may modify the existing argument about the order of the chiral phase transition in the $N_f = 2$ case.

The properties of quarks and gluons at high temperature are key ingredients for understanding the evolution of the Universe and the heavy ion collision experiment. Lattice QCD is the most reliable formulation of QCD for the investigation of non-perturbative properties of quarks and gluons, and indeed a lot of works from the early stage of lattice gauge theories were devoted to clarify them [1]. Although many interesting and useful results were obtained, still there remain unsolved issues. Among other things, it is a long standing issue to determine the order of the chiral transition for $N_f = 2$ QCD, which has recently attracted a lot of attention [2].

In this article we give a new perspective on the properties of quark-gluon state above the chiral phase transition temperature T_c, and thereby give a new insight into the order of the chiral phase transition. Our main idea is to use the concept of the conformal field theory with an IR cutoff. After presenting the renormalization group argument and conjecturing that the propagator decays with a power-law corrected Yukawa-type decaying form, we discuss numerical results of lattice calculations and their physical implications on the chiral transition in $N_f = 2$ QCD. Some preliminary results have been presented in [3].

We discuss lattice QCD at high temperature for small N_f ($2 \leq N_f \leq 6$) massless fermions in fundamental representation where the chiral phase transition occurs at some critical temperature T_c. Our general argument that follows can be applied to any number of flavors ($2 \leq N_f \leq 6$) with any formulation of gauge theories on the lattice.

For numerical simulations in this article we take $N_f = 2$ and employ the Wilson quark action and the standard one-plaquette gauge action on the Euclidean lattice of the size $N_x = N_y = N_z = N$ and N_t, with the lattice spacing a. We impose an anti-periodic boundary condition in the time direction for fermion fields and periodic boundary conditions otherwise. In order to obtain physical quantities at temperature T, we have to take the thermodynamic limit $N \to \infty$ and the continuum limit $a \to 0$, keeping $N_t\, a = 1/T$ fixed. The order of the thermodynamic limit and the continuum limit can be exchanged.

The theory is defined by two parameters; the bare coupling constant g_0 and the bare degenerate quark mass m_0 at ultraviolet (UV) cutoff. We also use, instead of g_0 and m_0, $\beta = 6/g_0^2$ and $K = 1/2(m_0 a + 4)$. We define the quark mass m_q through Ward-Takahashi identities [4, 5] with renormalization constants being suppressed.

We first consider the case that the renormalized quark mass is zero. Then the renormalized coupling constant is the only relevant variable in the theory. A running coupling constant $g(\mu; T)$ at temperature T can be defined as in the case of $T = 0$. The following discussion can be applied to any definition of the running coupling constant $g(\mu; T)$. One promising way is the Wilson MCRG method in order to investigate the infrared behavior.

In most of the existing literatures (see e.g. ref. [6]), the authors discussed the running coupling constant $g(\mu; T)$ mainly in the UV regime. However, what we would like to understand is the IR behavior in connection with the issues raised in the introduction.

In the UV regime, since the theory is asymptotically free, the running coupling constant at finite T can be expressed as a power series of the running coupling constant at $T = 0$ as long as g is small [7]. The leading term is universal in the limit $g \to 0$.

However, in the IR region, $g(\mu; T)$ is quite different from $g(\mu; T = 0)$, since the IR cutoff $\Lambda_{\rm IR}$ in the time direction is T, while the IR cutoff is zero at zero temperature. Furthermore, when $T/T_c > 1$, where the quark is not confined, the running coupling constant $g(\mu; T)$ cannot be arbitrarily large. This means that there is an IR fixed point with non-trivial zero of the beta function when $T/T_c > 1$. This is the key observation in this article.

As long as $T < T_c$, the beta function is negative all through g. As the temperature is increased further, the form of the beta function will change as in Fig.1: (left) When $T > T_c$ but $T \sim T_c$, the beta function changes the sign from negative to positive at large g; (middle)



FIG. 1. The beta function $\beta(g(\mu;T))$ at finite temperature T

As temperature increases the fixed point moves toward smaller g; (right) When $T \gg T_c$ it changes the sign at small g.

In Lorentz invariant theories, the existence of an IR fixed point implies that the theory is conformal (see e.g. [8] for a review). In our case with finite temperature, we argue that the consequence of the IR fixed point for $T > T_c$ naturally leads to the conformal field theories with an IR cutoff we have recently investigated in [10]. There, the dynamics of conformal theories was realized in the SU(N) gauge theory, when the number of flavors N_f is within the conformal window [9], and the finite lattice size introduced the IR cutoff. The discussion presented there can be applied here with small modifications. We refer ref. [10] for the RG argument which is a background of the predictions made in ref. [10] and here.

To verify the above idea of conformal field theories with an IR cutoff realized in high temperature QCD, we will study the propagator of a local meson operator $G_H(t)$ on a finite lattice for $T > T_c$ with T_c defined for the chiral transition temperature in $m_q \to 0$ limit. The analogy to the conformal field theories with an IR cutoff defined for the large N_f (within the conformal window) QCD implies that the behavior of $G_H(t)$ qualitatively differs depending on whether the quark mass is smaller than the critical mass or not.

When the theory is in the relatively heavy quark region, it decays exponentially at large t as

$$G_H(t) = c_H \, \exp(-m_H t), \tag{1}$$

where m_H is the energy of the ground state in the channel H. In contrast to the "confining phase" m_H is not necessarily the mass of a meson. It corresponds to the eigenvalue of the ground state energy in the channel H.

The conjecture made in ref. [10] which was confirmed there can be translated to the following conjecture: When $T/T_c > 1$, in the "conformal region" defined by

$$m_H \le c\,\Lambda_{\rm IR} \tag{2}$$

where c is a constant of order 1, the propagator $G_H(t)$ behaves at large t as

$$G_H(t) = \tilde{c}_H \frac{\exp(-\tilde{m}_H t)}{t^{\alpha_H}}, \tag{3}$$

which is a power-law corrected Yukawa-type decaying form instead of the exponential decaying form observed when $m_H > c\,\Lambda_{\rm IR}$.

Now let us discuss the results of our numerical calculations in the $N_f = 2$ case on a $16^3 \times 64$ lattice for several sets of parameters. The algorithm we employ is the blocked HMC algorithm [12].

After confirming the chiral transition occurs around $K = 0.151$ at $\beta = 6.0$, we choose the following values of $\beta' s$: $\beta = 6.5$ ($T \sim 2\,T_c$), $\beta = 7.0$ ($T \sim 4\,T_c$), $\beta = 8.0$ ($T \sim 16\,T_c$), $\beta = 10.0$ ($T \sim 100\,T_c$), and $\beta = 15.0$ ($T \sim 10^5\,T_c$). We take several values of K for each β in such a way that the quark masses $m_q' s$ take values $0.00 \le m_q \le 0.10$. The crude estimates of the temperature are based on $\Delta\beta \sim 0.5$ for the scale change of a factor 2 in the lowest perturbation.

We choose run-parameters in such a way that the acceptance of the global metropolis test is about 70%. The statistics are 1,000 MD trajectories for thermalization and 1,000 MD trajectories or 500 MD trajectories for the measurement. We estimate the errors by the jack-knife method with a bin size corresponding to 100 HMC trajectories.

We define the effective mass $m_H(t)$ by

$$\frac{\cosh(m_H(t)(t - N_t/2))}{\cosh(m_H(t)(t + 1 - N_t/2))} = \frac{G_H(t)}{G_H(t+1)}$$

.

FIG. 2 shows the t dependence of the effective mass for the PS channel in the case $T \simeq 100 T_c$ with three types of sources. On the left panel, we see the clear plateau of the effective mass at $t = 22 \sim 31$ when quark mass is relatively large; $m_q = 0.30$ ($K = 0.125$). On the other hand, on the right panel, we see the effective mass is slowly decreasing without no plateau up to $t = 31$ when the quark mass is small; $m_q = 0.028$ ($K = 0.135$), suggesting the power-law correction. The power-law corrected fit for the local-local data in the $K = 0.135$ case with the fitting range $t = [15:31]$ with $\alpha_H = 1.1(1)$ reproduces the data very well.

We show the effective mass plot for the local source - local sink in two more cases for $T \simeq 4T_c$ and $T \simeq 10^5\,T_c$ in Fig.(3), together with the power-law corrected fit with the fitting range $t = [15:31]$. Both of the fit for $\beta = 7.0$ at $K = 0.142$ with $\alpha_H = 0.7(1)$ and the fit for $\beta = 15.0$ at $K = 0.130$ with $\alpha_H = 1.21(4)$ reproduce the data for $t = [15:31]$ very well.

For all cases with $0.0 \le m_q \le 0.10$, we observe the similar power corrected behavior as shown in Fig. 2 and we confirm the fit similar to those shown in Fig.



FIG. 2. The effective mass for $N_f = 2$ at $\beta = 10.0$ (left: for $K = 0.125$; exponential decay) and (right: for $K = 0.135$; power-law corrected Yukawa decay): With three types of sources; the local-sink local-source (black squares), local-sink doubly-smeared-source (red circles) and local-sink doubly-wall-source (green triangles)



FIG. 3. The effective mass plots for local-sink local-source case and fits by power-law corrected Yukawa type decay (left: for $\beta = 7.0$ at $K = 0.142$) and (right: for $\beta = 15$ at $K = 0.130$)

3 well reproduces the data. Thus we have verified our conjecture that the propagator on a finite lattice in the conformal region given by eq.(2) behaves at large t as a power-law corrected Yukawa-type decaying form given by eq.(3). We have not systematically estimated the constant c in eq.(2). In general, the c depends on N_f and β. We quote $c \simeq 2.0 \times 2\pi$ at $\beta = 6.0$ for $N_f = 7$ and $\beta = 11.5$ for $N_f = 16$ [10] for later use.

Our numerical simulation confirms that the behavior of $G_H(t)$ is similar to each other between the small N_f QCD at high temperature and the large N_f QCD with an IR cutoff. One reason behind is that once gauge coupling constant is at the IR fixed point, the renormalization group equation is characterized by only one number, the mass anomalous dimension which is related to α_H. All the results show the existence of the "conformal region" for a wide region for $T > T_c$ on a finite lattice. We believe it will survive in the thermodynamic limit from our theoretical argument. We hope to revisit more rigorous treatment of the thermodynamic limit in the future work.

In relation to the thermodynamic limit, we would like to point out that there is one difference between the small N_f QCD at high temperature and the large N_f QCD with an IR cutoff in the lattice size dependence because the origin of the IR cutoff is quite different. To understand this point, let us consider the phase diagram in terms of β and K for the small $N_f (N_f = 2 \sim 6)$ QCD on a finite lattice. The chiral phase transition point on the massless quark line moves towards larger β with larger lattice, since the beta function is non-positive all through β at $T = T_c$. (For $N_f = 2$ the chiral transition point $\beta_{ct} = 3.9 \sim 4.0$ [11] at $N_t = 4$ moves to $\beta_{ct} \simeq 6.0$ at $N_t = 64$. For $N_f = 6$ $\beta_{ct} \simeq 0.3(1)$ [11] at $N_t = 4$ and the movement of β_{ct} is very slow, which might be due to the smallness of the beta function as is in the walking technicolor theory.) Assume we are in the deconfining phase at some β with some lattice size. When the lattice size is increased with β fixed, the deconfining phase changes to the confining phase at some lattice size. This is clearly different from the large N_f QCD within the conformal window in which the IR fixed point remains even in the infinite volume limit at zero temperature.

Now we would like to discuss the implication of the existence of the IR fixed point in high temperature QCD and the existence of the "conformal region" on a finite lattice. In this article we concentrate on the issue of the order of the chiral phase transition in the $N_f = 2$ case. Besides our numerical results, our key observation is the existence of an IR fixed point at $T > T_c$. We stress that the reasoning for the existence can be justified even in the thermodynamic limit.

Pisarski and Wilczek [13] mapped $N_f = 2$ QCD at high temperature to the three dimensional sigma model and pointed out that if $U_A(1)$ symmetry is not recovered at the chiral transition temperature, the chiral phase transition of QCD in the $N_f = 2$ case is 2nd order with exponents of the three dimensional $O(4)$ sigma model.

The $O(4)$ scaling relation was first tested for staggered quarks [14]. For the Wilson quarks it was shown that the chiral condensate satisfies remarkably the $O(4)$ scaling relation, with the RG improved gauge action and the Wilson quark action [15] and with the same gauge action and the clover-improved Wilson quark action [16] (See, for example, Fig.6 in ref. [15]). It was also shown for staggered quarks the scaling relation is satisfied in the $N_f = 2+1$ case [17], extending the region from $T/T_c > 1$ adopted in [15] and [16] to the region including $T/T_c < 1$. These results imply the transition is second order.

However, recently, it was shown that the expectation value of the chiral susceptibility $\chi_\pi - \chi_\sigma$ is zero [18] in thermodynamic limit when the $SU(2)$ chiral symmetry is recovered under the assumptions we will discuss below. This is consistent with that the $U_A(1)$ symmetry is recovered, which implies the transition is 1st order according to [13]. Apparently the two conclusions are in contradiction.

Here we revisit this issue with the new insight of conformal field theories with an IR cutoff. We have to check two points: (1) whether the mass region investigated belongs to the conformal region; (2) whether the power-corrected behavior of the propagators is properly taken into account. The chiral transition occurs at the massless quark line and therefore the mass region, where the $O(4)$ scaling holds, will be in the conformal region. We took the quark mass region $m_q \leq 0.4$ in the fit of Fig.6 of ref.[15]. On the other hand, the critical pion mass m_π for the boundary of the conformal region is estimated using the general equation $m_\pi = c\,\Lambda_{\rm IR}$ with $\Lambda_{\rm IR} = (N^3 \times N_t)^{1/4}$ as in [10] since the spatial lattice size is small 8^3 with $N_t = 4$. It gives $m_\pi \simeq 1.9$, with $c \simeq 2.0 \times 2\pi$. Since the quark mass region $m_q \leq 0.4$ corresponded to $m_\pi \leq 1.7$, we conclude that the mass region studied there belonged to the conformal region. This explains the fact that more than 30 data were excellently on the given smooth scaling curve with only one fitting parameter, the chiral transition coupling constant, besides the scales of x-axis and y-axis.

One more point is concerned with the power corrected Yukawa-type decay. We calculated the chiral condensate subtracted using [4]

$$\langle \bar\Psi\Psi \rangle_{\rm sub} = 2 m_q a Z \sum_{x,t} \langle \pi(x,t)\pi(0) \rangle,$$

where the sum is over space-time. This is equivalent to the time integration of the propagator with power corrected Yukawa-type decay. Note that this formula itself is always valid irrespective of the form of the propagator, but the scaling of the summed up quantity is affected by the power corrections. In [15], we have directly studied the chiral condensate without noticing the power corrected decaying form, but the sum automatically takes into account the effects. Thus the analysis there is consistent with the new insight of conformal field theories with an IR cutoff and therefore the conclusion does not change that the $O(4)$ scaling relation is remarkably satisfied and the phase transition must be 2nd order.

On the other hand, it is assumed in ref. [18] that the vacuum expectation value of mass-independent observable is an analytic function of m_q^2, if the chiral symmetry is restored. However, in the conformal region the propagator of a meson behaves as eq.(3) and the relation between the m_H and the m_q is given by the hyperscaling relation [19]

$$m_H = c\, m_q^{1/(1+\gamma*)},$$

with γ^* the anomalous mass dimension. This anomalous scaling implies m_H is not analytic in terms of m_q^2 and the analyticity assumption does not hold. It should be noted that the Ward -Takahashi Identities in [18] are proved in the thermodynamic limit and therefore the numerical verification of the hyperscaling in the limit will be decisive. We stress however that the hyperscaling is theoretically derived with the condition of the existence of the IR fixed point and multiplicative renormalization of m_q. We believe that this violation of the analyticity assumption resolves the apparent discrepancy as also mentioned in [18] as a viable possibility.

In the article in preparation we hope to discuss other physical implications of conformal field theories with an IR cutoff realized at high temperature QCD which are not discussed in this article, together with the discussion on the relation between large N_f QCD within the conformal window and small N_f QCD at high temperature.

ACKNOWLEDGMENTS

We would like to thank S. Aoki, H. Fukaya, K. Kanaya and Y. Taniguchi for useful discussion. The calculations were performed with HA-PACS computer at CCS, University of Tsukuba and SR16000 at KEK. We would like to thank members of CCS and KEK for their strong support for this work.

---

[1] J. Kuti, J. Polonyi, and K. Schachlanyj, Phys. Lett. 98B (1981) 199; J. Engels, F. Karsch, H. Satz, I. Montvay, Phys. Lett. B101 (1981) 89 . L. McLerran and B. Svetitsky, Phys. Rev. D24 (1981) 450;

[2] M. P. Lombardo, arXiv:1301.7324]hep-lat] and references quoted.

[3] Y. Iwasaki, to be published in Proceedings of SCGT 12 held in Nagoya on December 4 -7,2012: arXiv:1212.4343.

[4] M. Bochicchio et al., Nucl. Phys. **B262** (1985) 331.

[5] S. Itoh, Y. Iwasaki, Y. Oyanagi and T. Yoshié, Nucl. Phys. **B274** (1986) 33.

[6] O. Kaczmarek, F. Karsch, F. Zantow and P. Petreczky, Phys. Rev. **D70**(2004) 074505.

[7] M. Luescher, P. Weisz and U. Wolff Nucl. Phys. B359 (1991) 221.

[8] Y. Nakayama, arXiv:1302.0884 [hep-th].

[9] T. Banks and A. Zaks, Nucl. Phys. **B196** (1982) 189.

[10] K.-I. Ishikawa, Y. Iwasaki, Yu Nakayama and T. Yoshie, Phys. Rev. D (to be published); arXiv:1301.4785.

[11] Y. Iwasaki, K. Kanaya, S. Kaya, S. Sakai, and T. Yoshié, Phys. Rev. **D54** (1996) 7010.

[12] M. Hayakawa, et al., PoS LATTICE **2010** (2010) 325.

[13] R, Pisarski and F. Wilczek, Phys. Rev. **29** (1984) 338.

[14] F. Karsch, Phys. Rev. D49 (1994) 3791; F. Karsch and E. Laermann, ibid. D50 (1994) 6954.

[15] Y. Iwasaki, K. Kanaya, S. Kaya, and T. Yoshié, Phys. Rev. Lett. **78** (1997) 179.

[16] A. Ali khan et al. (CP-PACS Collaboration), Phys. Rev. **D63** (2001) 034502.

[17] S. Ejiri et al., Phys. Rev. **D80**(2009) 094505; A. Bazavov et al., Phys. Rev. **D85**(2012) 054503.

[18] S. Aoki, H. Fukaya, Y. Taniguchi, arXiv:1209.2061

[19] V.A. Miransky, Phys. Rev. **D59** (1999) 105003; L. Del Debbio and R. Zwicky, Phys. Rev. **D82** 014502.